

ISO AND THE COSMIC INFRARED BACKGROUND

Hervé Dole¹¹ SIRTf/MIPS Team, Steward Observatory, University of Arizona, 933 N Cherry Ave, Tucson, AZ, 85721, USA
http://lully.as.arizona.edu

ABSTRACT

ISO observed, for the first time to such a high sensitivity level, the mid- and far- infrared universe. A Number of deep surveys were performed to probe the cosmological evolution of galaxies. In this review, I discuss and summarize results of mid-infrared ISOCAM and far-infrared ISOPHOT surveys, and show how our vision of the extragalactic infrared universe has become more accurate. In particular, ISO allowed us to resolve into sources a significant fraction of the Cosmic Infrared Background (CIB) in the mid-infrared, and to probe a fainter population in the far-infrared with the detection of the CIB fluctuations. Together with other wavelength data sets, the nature of ISO galaxies is now in the process of being understood.

I also show that the high quality of the ISO data put strong constraints on the scenarios of galaxy evolution. This induced a burst in the development of models, yielding to a more coherent picture of galaxy evolution.

I finally emphasize the potential of the ISO data archive in the field of observational cosmology, and describe the next steps, in particular the forthcoming cosmological surveys to be carried out by SIRTf.

Key words: ISO – infrared galaxies – galaxy evolution – extragalactic surveys – cosmic infrared background

1. INTRODUCTION

The Cosmic Infrared Background (CIB) (e.g. Puget et al, 1996; Hauser & Dwek, 2001) is the relic emission of galaxy formation and evolution, being composed of light radiated from galaxies since their formation. The aim of the cosmological surveys is to resolve the CIB into sources, and to provide data on galaxies at various wavelengths, redshifts, with enough quality on individual objects as well as with statistical significant samples, in order to address the questions of the processes of galaxy formation and evolution, and the nature of the galaxies.

The CIB Spectral Energy Distribution (SED), shown in Fig. 1 (from Hauser & Dwek, 2001) shows the existence of a minimum between 3 and 10 μm separating direct stellar radiation from the FIR part due to radiation re-emitted by dust. The latter radiation contains at least a comparable integrated power as the former, and perhaps as much




Figure 1. The Cosmic Infrared Background Spectral Energy Distribution from Hauser & Dwek (2001). This plot includes detection (solid line in the submm part), upper limits (triangles and diamonds), and lower limits from source counts (squares). The shaded region represents current observational limits, and the solid rising curve in the mm range represents the CMB..

as 2.5 times more. This ratio is much larger than what is measured locally ($\sim 30\%$). The CIB is thus likely to be dominated by a population of strongly evolving redshifted IR galaxies.

ISO revolutionized our view of the extragalactic universe with sensitive cosmological surveys carried out in the mid- and far- infrared. Of course, other cosmological surveys have been carried out with SCUBA, MAMBO, Chandra, XMM, and in the optical range; all give invaluable informations about galaxy evolution, but it is beyond the scope of this short paper to review them. Reviews of extragalactic results from ISO can be found in Genzel & Cesarsky (2000) and Franceschini et al. (2001), and a comprehensive overview of ISO operations and science in Casoli et al. (2000) (particularly in the Puget & Guiderdoni paper). I will here summarize the main results obtained in the MIR with ISOCAM (Sect. 2), in the FIR with ISOPHOT (Sect. 3), and overview the models that ISO data were able to constrain (Sect. 4). Sect. 5 emphasizes the potential of the ISO archive, and in Sect. 6 I




Figure 2. *K*-corrections from Franceschini et al (2001). Dash: M82 spectrum at $7\ \mu\text{m}$. Dot: inactive spiral at $15\ \mu\text{m}$. Line: M82-like spectrum at $15\ \mu\text{m}$.

briefly explore the future of IR (and mm) astronomy, in particular with SIRTf.

2. MID INFRARED SURVEYS

2.1. SPECIFICITIES

The mid-infrared surveys have been conducted with ISO-CAM (Cesarsky et al., 1996), mainly at 7 and $15\ \mu\text{m}$; a few have been performed at $4.5\ \mu\text{m}$ and $12\ \mu\text{m}$ (Clements et al., 1999). At $7\ \mu\text{m}$ using the LW2 filter (e.g. Taniguchi et al., 1997; Oliver et al., 2002), the contamination by stars is important and data at other wavelengths (usually near infrared) are needed to identify them. The decreasing *k*-correction is less favorable for detecting higher redshift galaxies (dash line in Fig. 2, taken from Franceschini et al., 2001).

At $15\ \mu\text{m}$ using the LW3 filter (e.g. Elbaz et al., 1999), the stellar contamination is less of a problem and the *k*-correction of galaxies is favorable for detecting sources up to redshifts around 1.4 (Fig. 2). Thus, a significant part of the extragalactic results are based on $15\ \mu\text{m}$ observations.

2.2. SOURCE COUNTS

Source counts at $15\ \mu\text{m}$ show an impressive consistency over four decades of flux between surveys of different depths (e.g. Elbaz et al., 1999, Gruppioni et al, 2002). Data come from large and shallow surveys like ELAIS (e.g. Serjeant et al, 2000), to narrower and deeper surveys (e.g. Serjeant et al, 1997, Aussel et al, 1999; Altieri et al, 1999). The number counts (summarized in Fig. 3 by Elbaz et al.,




Figure 3. Differential source counts (normalized to Euclidean) at $15\ \mu\text{m}$ from Elbaz et al (1999). The shadowed area represents a scenario with no evolution. Symbols represent different surveys.

1999) are in excess by a factor of 10, at the 0.5 mJy level, compared to non evolution scenarios. The slope is steep ($\alpha = 3.0 \pm 0.1$ in the differential counts, for $S_\nu > 0.4$ mJy), and a turnover appears at $S_\nu = 0.4$ mJy. At fainter levels, the convergence (decrease) is fast.

2.3. NATURE OF THE GALAXIES

With a median redshift of $z \sim 0.8$ and most of the sources with $0.5 < z < 2$ (Aussel et al., 1999), the MIR sources have a mean luminosity of $6 \times 10^{11} L_\odot$ (Elbaz et al., 2002). Most of them are experiencing intense stellar formation of about $100 M_\odot \text{yr}^{-1}$, which appears to be uncorrelated with the faint blue galaxy population dominating the optical counts at $z \sim 0.7$ (Ellis 1997; Elbaz et al. 1999). Fig. 4 (from Elbaz et al., 2002) represents their HDF-N sample, and summarizes the properties of MIR galaxies: redshift distribution, luminosity, and SFR (for 80% of the non-AGN galaxies).

2.4. GALAXY EVOLUTION

Elbaz et al (2002) present a summary of the galaxy evolution at $15\ \mu\text{m}$ and beyond. First, they show that 60% of the CIB at $15\ \mu\text{m}$ is created by LIRGs ($L > 10^{11} L_\odot$). Second, they show that the comoving luminosity density at $15\ \mu\text{m}$ was about 55 times larger at $z \sim 1$ than today. Third, still using some assumption about the SED of the galaxies, they conclude that the comoving density of IR luminosity radiated by dusty starbursts (which is directly related to the star formation) was about 70 ± 35 times larger at $z \sim 1$ than today. At intermediate redshifts, Flores et al (1999) also determined the star formation rate.




Figure 4. Infrared luminosity (left axis) and star formation rate (right axis, for starbursts only) of $15 \mu\text{m}$ $S > 0.1$ mJy galaxies in the HDF-N from Elbaz et al (2002). The dash curve represents the sensitivity limit. This figure also illustrates the redshift distribution of the sources, with a median redshift of $z = 0.8$ (Aussel et al., 1999).

Finally, they show that about half of the CIB at $140 \mu\text{m}$ is produced by LIRGs and about one third by ULIRGs. Thus MIR galaxies observed by ISOCAM have resolved about 75% of the CIB at $140 \mu\text{m}$.

3. FAR INFRARED SURVEYS

3.1. SPECIFICITIES

The far-infrared surveys have been conducted with ISOPHOT (Lemke et al., 1996), mainly at 90 and $170 \mu\text{m}$; a few have been performed at 60, 120, 150 and $180 \mu\text{m}$ (Juvela et al., 2000; Linden-Voernle et al., 2000). In the 50 to $100 \mu\text{m}$ range, the C100 Ge:Ga camera is sensitive to the peak of rest-frame emission from obscured star formation. The decreasing k-correction, in addition to a challenging data processing, led to probe mainly the local universe. With 46 arcseconds per pixel and a FWHM of about the same size, the angular resolution at $100 \mu\text{m}$ is significant improved since IRAS.

In the 100 to $200 \mu\text{m}$ range, the C200 stressed Ge:Ga camera is sensitive to the very cool local galaxies as well as higher redshift starburst galaxies, thanks to an advantageous k-correction (e.g. Guiderdoni et al., 1997) and the behavior of the camera (e.g. Lagache & Dole, 2001). With 92 arcseconds per pixel and a FWHM of about the same size, the angular resolution at $170 \mu\text{m}$ is an issue for source identifications.

3.2. SOURCE COUNTS

Source counts at $170 \mu\text{m}$ (e.g. Kawara et al., 1998; Puget et al., 1999; Dole et al., 2001) exhibit a steep slope of




Figure 5. Distribution of IR luminosities from Patris et al (2002). Solid line: FIRBACK FSM $170 \mu\text{m}$ sample ($S > 200$ mJy); Dash: IRAS $100 \mu\text{m}$ sample from Bertin et al (1997) with $110 < f_{60} < 200$ mJy.

$\alpha = 3.3 \pm 0.6$ between 180 and 500 mJy (lower left panel on Fig. 7) and, like in the MIR range, show sources in excess by a factor of 10 compared with no evolution scenario. This strong evolution was unexpected and quite difficult to reproduce with models. Matsuhara et al (2002) extended the source count analysis to fainter fluxes and still detected a strong evolution.

At 60 and $90 \mu\text{m}$, the situation is less clear, but preliminary source counts (Kawara et al., 1998; Efsthathiou et al., 2000; Linden-Voernle et al., 2000) are compatible with no evolution scenarios on their bright end, and begin to show evolutionary effects on their bright end. New processing techniques (see Sect. 5) will certainly allow to go deeper.

3.3. NATURE OF THE GALAXIES

Determining the nature of the FIR galaxies has been a longer process than in the MIR, mainly because of the difficulty to find the shorter wavelength counterparts in a large beam. Various techniques have been used to overcome this problem, one of the most successful being the identification using 20cm radio data (e.g. Ciliegi et al., 1999). Another technique is the FIR multiwavelength approach (Juvela et al., 2000) that helps constraining the position and the SED; it also helps to separate the cirrus structures from the extragalactic sources. A variation is to use ISOCAM and ISOPHOT data, like the ELAIS Survey (Oliver et al., 2000; Serjeant et al., 2000, 2001). Finally, the Serendipity Survey (Stickel et al., 1998, 2000), by covering large and shallow areas, allows to detect many bright objects easier to follow-up or already known.

FIR ISO galaxies can be sorted schematically into two populations. First, the low redshift sources, typically $z <$




Figure 6. Fluctuations of the CIB in a power spectrum analysis of the FIRBACK/ELAIS N2 field at 170 microns by Puget & Lagache (2000). Observed power spectrum: diamond; straight continuous line: the best fit cirrus power spectrum; dash line: cirrus power spectrum deduced from Miville-Deschênes et al (2002); continuous curve: detector noise.

0.3 (e.g. Serjeant et al., 2001; Patris et al., 2002; Kakazu et al., 2002), have moderate IR luminosities, below $10^{11} L_{\odot}$, and are cold (Stickel et al., 2000). Second, sources at higher redshift, $z \sim 0.3$ (Patris et al., 2002, see Fig. 5) and beyond, $z \sim 0.9$ (Chapman et al., 2002) are more luminous, typically $L > 10^{11} L_{\odot}$, and appear to be cold. Serjeant et al. (2001) derived the Luminosity Function at $90 \mu m$, and started to detect an evolution compared to the local IRAS $100 \mu m$ sample.

3.4. FLUCTUATIONS OF THE CIB

Sources below the detection limit of a survey create fluctuations. If the detection limit does not allow to resolve the sources dominating the CIB intensity, which is the case in the FIR with ISO, characterizing these fluctuations gives very interesting information on the spatial correlations of these unresolved sources of cosmological significance. An example of the modeled redshift distribution of the unresolved sources at $170 \mu m$ can be found in Fig. 12 of Lagache et al (2002); the sources dominating the CIB fluctuations have a redshift distribution peaking at $z \sim 0.9$. After the pioneering work of Herbstmeierer al (1998) with ISOPHOT, Lagache & Puget (2002) discovered them at $170 \mu m$ in the FIRBACK data, followed by other works at 170 and $90 \mu m$ (Matsuhara et al., 2000; Puget & Lagache, 2000; Kiss et al, 2001). Fig. 6 shows the CIB fluctuations in the FN2 field by Puget & Lagache, (2000), at wavenumbers $0.07 < k < 0.4 \text{ arcmin}^{-1}$.

4. MODELS

ISO provided high quality data, taken in the frame of the cosmological survey programs, as well as nearby galaxies or serendipitous programs. The first observable to be widely used as a constraint on the models is the source counts at 15 and $170 \mu m$, but also at 7 and $90 \mu m$. The redshift distributions, mainly at $15 \mu m$ (and soon at $170 \mu m$) put additional constraints, as well as a better knowledge of the nature of the galaxies: AGN vs Starburst, luminosities, star formation rate. The global star formation rate, as derived in part from ISOCAM data, adds other constraints to the models. Finally, a few models use explicitly the constraint of the level of the CIB fluctuations in the far infrared.

These observables and constraints induced a burst in the development of new models in the late nineties, among which: Roche & Eales (1999), Tan et al. (1999), Devriendt & Guiderdoni (2000), Dole et al. (2000), Wang (2000), Chary & Elbaz (2001), Franceschini et al. (2001), Malkan & Stecker (2001), Pearson (2001), Rowan-Robinson (2001), Takeuchi (2001), Xu et al. (2001), Balland et al. (2002), Lagache et al. (2002), Totani & Takeuchi (2002), Wang (2002). This development was also made possible because of the better knowledge of the galaxies SED in the infrared (e.g. Dale et al., 2000; Helou et al., 2000; Lutz et al., 2000; Stickel et al., 2000, Tuffs & Popescu, 2002), with the availability of the SED of the CIB (Gispert et al., 2000; Hauser & Dwek, 2001), and of complementary data sets (X-rays, optical, NIR, submm, radio).

Even if there is no unique scenario reproducing all the observables, the models help now to have a more coherent picture of galaxy evolution. It is beyond the scope of this paper to review the models and their predictions, we thus give here only two examples. Fig. 7 shows the observed source counts at 15 , 60 , 170 and $850 \mu m$ and the fits by Lagache et al (2002). Using other kind but similar evolutions (pure density and pure luminosity evolutions in addition to the density+luminosity evolution), Chary & Elbaz (2001) predict the star formation rate and compare it with data and Xu et al (2001) (Fig. 8). Combined with the different approach of Gispert et al (2000), the SFR is well constrained in the redshift range $1 < z < 3$.

5. POTENTIAL OF ISO DATA

Even if the ISO data led to an impressive scientific return with a high efficiency (about 62% of the data are published) in the field of the cosmological evolution of the galaxies, many other results are still to come, based on the amount of unpublished data available in the archive.

In addition, new techniques of data reduction are now available, thanks to a better understanding of detector behaviors, both for the Silicon Array (CAM)(Coulais & Abergel, 2000; Miville-Deschênes et al., 2000; Lari et al., 2001, 2002; Vaccari, 2002) and the Germanium detectors




Figure 7. Observed source counts at 15, 60, 170 and 850 μm with the model of Lagache et al (2002). Symbols represent data (CAM at 15 μm , IRAS at 60 μm , PHOT at 170 μm and SCUBA at 850 μm). Solid line: model prediction. Dash: LIRG/starburst component. Dot-dash: normal/cold galaxy component.

(PHOT) (Coulais et al., 2000). Processing unpublished data and/or reprocessing published data allows us to reach better sensitivity levels with a higher confidence (e.g. for ISOCAM HDF-N: Serjeant et al., 1997; Aussel et al., 1999; Désert et al., 1999. E.g. for ISOPHOT Lockman Hole: Kawara et al., 1998; Rodighiero et al., 2002).

Concerning the CIB and the cirrus foreground, a brilliant example of what can be done from the archive is the work of Kiss et al (2001,2002), detailed in this conference. Furthermore, the ISO data may induce discoveries in other data sets. A recent example concerns the fluctuations of the CIB in the far infrared. Lagache & Puget (2000) and Puget & Lagache (2000) measured the level of the fluctuations of the CIB at 170 μm in the FIRBACK fields. Given the SED of the CIB, it was possible to predict the fluctuation level at 60 and 100 μm (e.g. Lagache, Dole, Puget, 2002), and show that this might be detectable in the IRAS data. As a consequence, Miville-Deschênes et al. (2002) recently reported the first detection of the CIB fluctuations at 60 and 100 μm in the IRAS data.

6. THE NEXT STEPS: SIRTf AND BEYOND

SIRTf, the Space Infrared Telescope Facility, to be launched in early 2003, is expected to improve our view of the IR universe over the next five years, thanks to the recent technological developments in IR detector arrays. A number of multiwavelength cosmological surveys are scheduled with MIPS (Multiband Imaging Photometer for SIRTf) at 24, 70 and 160 μm (Dole et al., 2003¹) and IRAC (In-




Figure 8. Star Formation Rate from Chary & Elbaz (2001). Upper panel: min and max range from their model, and observed UV/opt data; dots represent Xu et al (2001) model. Lower panel: 3 different evolution scenarios from their model and data corrected for extinction. Line: pure luminosity; upper dash: pure density; lower dash: luminosity+density.

frared Array Camera) at 3.6, 4.5, 5.8 and 8.0 μm . In the frame of the Guaranteed Time (GTO) or Legacy (GOODS and SWIRE) Surveys, about 80 Sq. Deg. will be observed at different depths. Probing the properties of the galaxies with samples of statistical significance up to redshifts of 2.5 or more will be possible. Fig. 9 (from Dole et al, 2003) shows predictions of the redshift distribution at 24 μm for MIPS surveys.

The next steps after ISO and SIRTf for understanding the infrared and submillimeter Universe from space will be ISAS's ASTRO-F in 2004, ESA's Herschel & Planck (around 2007), and NASA's NGST (around 2010+), in addition to the ground-based ALMA (from 2005). These telescopes are the promise for a lot of exciting science to be done (at least) in the next 15 years !

ACKNOWLEDGEMENTS

I warmly acknowledge the organizers for having invited me to

¹ see also <http://lully.as.arizona.edu>




Figure 9. Predicted redshift distribution of sources observed at 24 μm with MIPS on SIRTf, from Dole et al (2003), for various 5σ surveys depths. Line: GTO shallow survey or SWIRE. Dash: GTO deep. Dot: GTO ultra-deep or GOODS.

give this review at this very interesting and enjoyable conference (especially for the tapas, the hot chocolate around the San Juan bonfire, and the cultural wonders). I also acknowledge the Programme National de Cosmologie and the Centre National d'Etudes Spatiales (CNES) for travel support for the work with my collaborators in France, and the funding from the MIPS project, which is supported by NASA through the Jet Propulsion Laboratory, subcontract # P435236.

REFERENCES

- Altieri B., Metcalfe L., Kneib J-P. et al., 1999, A&A, 343, L65
 Aussel H., Cesarsky C.J., Elbaz D., et al., 1999, A&A, 342, 313
 Balland C., Devriendt, J., Silk J., 2002, MNRAS, Submitted, astro-ph/0210030
 Bertin E., Dennefeld, M. & Moshir, M., 1997, A&A, 323, 685
 Casoli F., Lequeux J., David F., Eds, 'Infrared Space Astronomy, Today and Tomorrow', Les Houches Summer School LXX, EDP Sciences, Springer-Verlag, 2000
 Cesarsky C., Abergel, A., Agnese P., et al., 1996, A&A, 315, L32
 Chapman, S. C., Smail I., Ivison R., et al., 2002, ApJ, 573, 66
 Chary R., & Elbaz D., 2001, ApJ, 556, 562
 Ciliegi P., McMahon R. G., Miley G., et al., 1999, MNRAS, 302, 222
 Clements D., Desert F-X., Franceschini A., et al., 1999, A&A, 346, 383
 Coulais A., & Abergel A., 2000, A&AS, 141, 533
 Coulais A., Fouks B. I., Giovannelli J. F., et al., 2000, A&A, SPIE
 Dale D., Silbermann, N., Helou, G., et al., 2000, AJ, 120, 583
 Désert F-X., Puget J-L., Clements D., et al., 1999, A&A, 342, 363
 Devriendt J. & Guiderdoni B., 2000, A&A, 363, 851
 Dole H., Gispert R., Lagache, G., et al., 2000, Springer Lect. Notes in Ph., v548, 54, astro-ph/0002283
 Dole H., Gispert R., Lagache, G., et al., 2001, A&A, 372, 364
 Dole H., Lagache G., Puget J-L., 2003, ApJ, in press, astro-ph/0211312
 Efstathiou A., Oliver S., Rowan-Robinson M., et al., 2000, MNRAS, 319, 1169
 Elbaz D., Cesarsky C., Fadda D., et al., 1999, A&A, 351, L37
 Elbaz D., Flores H., Chanial P., et al., 2002, A&A, 384, 848
 Ellis R. S., 1997, ARAA, 35, 389
 Flores H., Hammer F., Thuan T. X., et al., 1999, ApJ, 517, 148
 Franceschini A., Aussel H., Cesarsky C., et al., 2001, A&A, 378, 1
 Genzel R., & Cesarsky C., 2000, ARAA, 38, 761
 Gispert R., Lagache G., Puget J-L., 2000, A&A, 360, 1
 Gruppioni C., et al, 2002, MNRAS, 335, 831
 Guiderdoni B., Bouchet F. R., Puget J-L., et al., 1997, Nature, 390, 257
 Hauser M., & Dwek E., 2001, ARAA, 37, 249
 Helou G., Lu N. Y., Werner M. W., et al., 2000, ApJ, 532, L21
 Herbstmeier U., Abraham P., Lemke D., et al., 1998, A&A, 332, 739
 Juvela M., Mattila K., Lemke D., 2000, A&A, 360, 813
 Kakazu Y., Sanders D., Joseph R., et al., 2002, A&A, IAU184, astro-ph/0201326
 Kawara K., Sato Y., Matsuhara H., et al., 1998, A&A, 336, L9
 Kiss C., Abraham P., Klaas, U., et al., 2001, A&A, 379, 1161
 Kiss C., et al., 2002, this conference
 Lagache G. & Puget J-L., 2000, A&A, 355, 17
 Lagache G. & Dole H., 2001, A&A, 372, 702
 Lagache G., Dole H., Puget J-L., 2002, A&A, MNRAS, in press, astro-ph/0209115
 Lari C., Pozzi F., Gruppioni C., et al., 2001, A&A, 325, 1173
 Lari C., et al., 2002, this conference
 Lemke D., Klaas U., Abolins J., et al., 1996, A&A, 315, L64
 Linden-Voernle M. J. D., Norgaard-Nielsen H. U., Jorgensenet H. E., et al., 2000, A&A, 359, L51
 Lutz D., 2000, New As. Rev., 44, L241
 Malkan M., & Stecker F., 2001, ApJ, 555, 641
 Matsuhara H., Kawara K., Sato Y., et al., 2000, A&A, 361, 407
 Miville-Deschênes M-A., Boulanger F., Abergel A., et al., 2000, 146, 519
 Miville-Deschênes M-A., Lagache G., Puget J-L., 2002, 393, 749
 Oliver S., Rowan-Robinson, M., Alexander D., et al., 2000, MNRAS, 316, 749
 Oliver S., Mann R. G., Carballo R., et al., 2002, MNRAS, 332, 536
 Patris J., Dennefeld M., Lagache G., et al., 2002, A&A, submitted
 Pearson C., 2001, MNRAS, 325, 1511
 Puget J-L., Abergel A., Bernard J-P., et al., 1996, A&A, 308, L5
 Puget J-L., Lagache G., Clements D., et al., 1999, A&A, 345, 29
 Puget J-L., & Lagache G., 2000, A&A, IAU204, astro-ph/0101105
 Roche N., & Eales S., 1999, MNRAS, 307, 111
 Rodighiero G., et al., 2002, this conference
 Rowan-Robinson M., 2001a, New As. Rev., 45, 631
 Rowan-Robinson M., 2001b, ApJ, 549, 745
 Serjeant S., Eaton N., Oliver S., et al., 1997, MNRAS, 289, 457

- Serjeant S., Oliver S., Rowan-Robinson M., et al., 2000, MNRAS, 316, 768
- Serjeant S., Efstathiou A., Oliver S., et al., 2001, MNRAS, 322, 262
- Stickel M., Bogun S., Lemke D., et al., 1998, A&A, 336, 116
- Stickel M., Lemke D., Klaas U., et al., 2000, A&A, 359, 865
- Takeuchi T., Ishii T., Hirashita H., et al., 2001, PASJ, 53, 37
- Tan J., Silk J., Balland C., 1999, ApJ, 522, 579
- Taniguchi Y., Cowie L., Sato Y., et al., 1997, A&A, 328, L9
- Totani T., & Takeuchi T., 2002, ApJ, 570, 470
- Tuffs R., & Popescu C., 2002, this conference
- Vaccari M., et al., 2002, this conference
- Xu C., Lonsdale C., Shupe D., et al., 2001, ApJ, 562, 179
- Wang Y., Biermann P., 2000, A&A, 356, 808
- Wang Y., 2002, A&A, 383, 755